

The Lattice NJL Model at Non-zero Baryon and Isospin Densities

David N. Walters^a and Simon Hands^{b*}

^aTheoretical Physics Group, Department of Physics and Astronomy, University of Manchester, Manchester M13 9PL, U.K.

^bDepartment of Physics, University of Wales Swansea, Singleton Park, Swansea SA2 8PP, U.K.

We present initial results of a numerical investigation of the chiral symmetry restoring transition in the (3+1)-dimensional Nambu – Jona-Lasinio model with both non-zero baryon chemical potential ($\mu_B \neq 0$) and isospin chemical potential ($\mu_I \neq 0$). With non-zero isospin chemical potential, the model suffers from a sign problem. We proceed in two ways: (i) We perform “partially quenched” simulations in which μ_I is made non-zero only during the measurement of chiral observables; (ii) We perform full simulations with imaginary isospin chemical potential with the aim to analytically continue results to real μ_I .

1. Motivation

At asymptotically high baryon chemical potential (μ_B) and low temperature (T) – where QCD can be treated in a perturbative manner – the ground-state of quark matter is found to be that of a colour-superconductor (for a recent review see e.g. [1]). Unfortunately, the sign problem prevents us from using lattice QCD to determine the ground-state at the more moderate densities typical in the cores of compact (neutron) stars.

One way to proceed is to study model field theories such as the Nambu – Jona-Lasinio (NJL) model. This purely fermionic field theory, in which colour-neutral quarks interact via a four-point contact term, not only contains the same global symmetries as two flavour QCD, but can be simulated on the lattice even with $\mu_B \neq 0$. In [2] we show that the ground-state of the lattice model with $\mu_u = \mu_d = \mu_B$, i.e. with “up” and “down” quarks sharing a common Fermi surface, exhibits s-wave superfluidity via a standard BCS pairing between quarks of different flavours; i.e.

$$\langle ud \rangle \neq 0; \quad \Delta_{BCS} \neq 0. \quad (1)$$

Within the cores of compact stars, however, the Fermi momenta k_F^u and k_F^d are expected to be

separate. A simple argument based on that of [3] suggests that for a two flavour Fermi liquid of massless quarks and electrons with $\mu_B = 400\text{MeV}$ and both weak equilibrium ($\mu_d = \mu_u + \mu_e$) and charge neutrality ($2n_u/3 - n_d/3 - n_e = 0$) enforced, all the Fermi momenta of the system are determined:

$$\begin{aligned} k_F^u &= \mu_u = \mu_B - \mu_e/2 = 355.5\text{MeV}, \\ k_F^d &= \mu_d = \mu_B + \mu_e/2 = 444.5\text{MeV}, \\ k_F^e &= \mu_e = 89\text{MeV}. \end{aligned} \quad (2)$$

The effect of separating the Fermi surfaces of pairing quarks in QCD should be to make the colour-superconducting phase less energetically favourable. Introducing $\mu_I \propto (\mu_u - \mu_d) \neq 0$ could prove a good method, therefore, to investigate the stability of the superfluid phase.

2. The Model

The action of the lattice NJL model (with $a \rightarrow 1$) is given by

$$\begin{aligned} S_{NJL} &= \sum_{xy} \bar{\Psi}_x M[\Phi, \mu_B, \mu_I]_{xy} \Psi_y \\ &+ \frac{2}{g^2} \sum_{\tilde{x}} \text{Tr} \Phi_{\tilde{x}}^\dagger \Phi_{\tilde{x}}, \end{aligned} \quad (3)$$

where $\Psi \equiv (u, d)^{\text{tr}}$ is the $SU(2)$ doublet of staggered up and down quarks defined on lattice sites x and $\Phi \equiv \sigma + i\vec{\pi} \cdot \vec{\tau}$ is a matrix of bosonic auxiliary fields defined on dual sites $\tilde{x}$. The fermion

*PPARC Senior Research Fellow

kinetic matrix M_{xy} is defined in [2] and we choose the same bare parameters used therein.

One can separate the Fermi surface of up and down quarks by simultaneously setting baryon chemical potential $\mu_B \equiv (\mu_u + \mu_d)/2 \neq 0$ and isospin chemical potential $\mu_I \equiv (\mu_u - \mu_d)/2 \neq 0$. With $\mu_I = 0$, $\tau_2 M \tau_2 = M^*$, which is a sufficient condition to show that $\det M$ is both real and positive [4]. With $\mu_I \neq 0$ however, this is no longer true such that once again we are faced with the sign problem.

The fact that physically the two scales are ordered $\mu_I \ll \mu_B$ suggests that one may be able to apply techniques recently developed to study QCD with $\mu_B \ll T$ [5,6,7]. First, however, we present the results of a partially quenched calculation.

3. Partially Quenched μ_I

Whilst the primary motivation for investigating $\mu_I \neq 0$ is to study the superfluid phase which sets in at large μ_B , this requires one to introduce an explicit symmetry breaking parameter (j); it is currently not clear how to study $\mu_I \neq 0$ in the $j \rightarrow 0$ limit [2]. Instead, we choose to study the chiral symmetry restoring phase transition with the aim of controlling the systematics of introducing $\mu_I \neq 0$.

The first step we take is to perform a “partially quenched” calculation in which $\mu_I = 0$ when generating the background fields and is made non-zero only during the measurement of fermion observables. In particular, we measure the up and down quark condensates

$$\langle \bar{u}u \rangle, \langle \bar{d}d \rangle \equiv \frac{1}{V} \frac{\partial \ln \mathcal{Z}}{\partial m_{u,d}} = \frac{1}{2} \langle \text{tr}(\mathbb{1} \pm \tau_3) M^{-1} \rangle \quad (4)$$

as functions of μ_B for various μ_I on a 12^4 lattice. Some results are presented in Fig. 1.

The results agree qualitatively with those of mean-field studies of the model in which the introduction of a small but non-zero μ_I is seen to suppress the up quark condensate and enhance the down [8,9]. This can be understood by noting that with $\mu_I \neq 0$, μ_u (μ_d) reaches the critical chemical potential at a lower (higher) value of μ_B . Why $\langle \bar{u}u \rangle$ deviates from the $\mu_I = 0$ result more




Figure 1. $\langle \bar{u}u \rangle$ and $\langle \bar{d}d \rangle$ condensates for various μ_B and μ_I on a 12^4 lattice.

than $\langle \bar{d}d \rangle$ is not so clear, but may be an effect associated with $T > 0$ on a finite lattice [9].

4. Continuation from Imaginary μ_I

One recent method used to study QCD with small baryon chemical potential μ_B is to measure quantities at imaginary μ_B , where the measure of the path integral is real, and fit results to a truncated Taylor expansion in μ_B/T about $\mu_B = 0$. One can then analytically continue the results to real μ_B [7]. We propose to use a similar method by simulating at imaginary isospin chemical potential ($\tilde{\mu}_I$), where once again $\det M$ is real, and expanding observables in powers of $\tilde{\mu}_I/\mu_B$.

For our initial investigation we have measured the quark condensates and baryon and isospin number densities

$$n_{B,I} \equiv \frac{1}{2V} \frac{\partial \ln \mathcal{Z}}{\partial \mu_{B,I}} = \frac{1}{4} \langle \bar{u} \gamma_0 u \pm \bar{d} \gamma_0 d \rangle \quad (5)$$

on a 12^4 lattice at $\mu_B = 0.6$, which from Fig. 1 can be seen to be where the effect of having $\mu_I \neq 0$ is largest. In QCD, one can show that e.g. $\langle \bar{\psi} \psi \rangle$ expanded about $\mu_B = 0$ is analytic in μ_B^2 , such that for small imaginary μ_B the quantity remains real [6]. For our simulations, however, this is not the case, and measured quantities are, in general, complex. Therefore, we fit the data by the Taylor series

$$\begin{pmatrix} \langle \bar{u}u \rangle \\ \langle \bar{d}d \rangle \end{pmatrix} = \sum_{n=0}^{\infty} \begin{pmatrix} A_n \\ B_n \end{pmatrix} \left(\frac{\tilde{\mu}_I}{\mu_B} \right)^n \quad (6)$$




Figure 2. Real and imaginary parts of $\langle\bar{u}u\rangle$ and $\langle\bar{d}d\rangle$ as functions of imaginary μ_I with $\mu_B = 0.6$ on a 12^4 lattice.

and

$$\begin{pmatrix} n_B \\ n_I \end{pmatrix} = \sum_{n=0}^{\infty} \begin{pmatrix} C_n \\ D_n \end{pmatrix} \left(\frac{\tilde{\mu}_I}{\mu_B} \right)^n, \quad (7)$$

each truncated at some suitable point. We can then analytically continue to real μ_I using e.g.

$$\langle\bar{u}u\rangle = A_0 + A_1\mu_I - A_2\mu_I^2 - A_3\mu_I^3 + \dots \quad (8)$$

Figure 2 shows the real and imaginary parts of the condensates as functions of $\tilde{\mu}_I$ with fits to *constant + quadratic* and *linear only* forms respectively. The quality of the fits is very good, suggesting that our ansätze are correct. The real parts are in complete agreement whilst the imaginary parts are anti-correlated, which implies that the sum of the condensates $\langle\bar{\psi}\psi\rangle$ contains only the even terms of the expansion, whilst their difference $\langle\bar{\psi}\tau_3\psi\rangle$ contains only the odd terms. A similar effect is seen in the number densities, as n_B is found to be real and therefore even in $\tilde{\mu}_I/\mu_B$, whilst the isospin density is pure imaginary. For a good quality fit to n_I , however, we have to include the cubic term. These data and fits are plotted in Fig. 3.




Figure 3. Baryon and isospin density as functions of imaginary μ_I with $\mu_B = 0.6$ on a 12^4 lattice.

Whilst these results are only preliminary, we have shown that we can calculate the coefficients in (6) and (7) as functions of μ_B and in principle, reproduce reliable forms of the curves in Fig. 1. With this aim, we plan to repeat this exercise for various values of μ_B in both the chirally broken and restored phases on various lattice volumes. Also, whilst it is difficult to study the diquark condensate in the $j \rightarrow 0$ limit, it would be interesting to compare the response of $\langle ud \rangle$ to μ_I at fixed j to that of $\langle\bar{\psi}\psi\rangle$ at fixed mass.

REFERENCES

1. D. H. Rischke, Prog. Part. Nucl. Phys. **52**, 197 (2004).
2. S. J. Hands and D. N. Walters, Phys. Rev. **D69**, 076011 (2004).
3. M. G. Alford, J. Berges, and K. Rajagopal, Nucl. Phys. **B571**, 269 (2000).
4. S. J. Hands *et al.*, Eur. Phys. J. **C17**, 285 (2000).
5. Z. Fodor and S. D. Katz, Phys. Lett. **B534** 87 (2002); Z. Fodor and S. D. Katz, JHEP **0203** 014 (2002).
6. C. R. Allton *et al.*, Phys. Rev. **D66**, 074507 (2002).
7. P. de Forcrand and O. Philipsen, Nucl. Phys. **B642**, 290 (2002); M. D’Elia and M.-P. Lombardo, Phys. Rev. **D67**, 014505 (2003).
8. D. Toublan and J. B. Kogut, Phys. Lett. **B564**, 212 (2003).
9. A. Barducci, R. Casalbuoni, G. Pettini, and L. Ravagli, Phys. Rev. **D69**, 096004 (2004).